

## Seismicity of the upper lithosphere and its relationships with the crust in the Italian region

S. SOLARINO<sup>1</sup> AND R. CASSINIS<sup>2</sup>

<sup>1</sup> *Centro Nazionale Terremoti, Istituto Nazionale di Geofisica e Vulcanologia, clo Dipteris, Università di Genova, Italy*

<sup>2</sup> *Viale Lombardia 30, Milano, Italy*

(Received: June 09, 2006; accepted: March 14, 2007)

**ABSTRACT** In a recent paper, we compared the earthquake hypocenters, plotted according to updated catalogs, with the structure of the Earth's crust interpreted after the results of seismic exploration (mainly the Deep Seismic Soundings - DSS). The comparison was made along several cross-sections in the Alpine range, the Italian peninsula and the surrounding seas. The main conclusions of that analysis were that 1) the majority of the events is positioned in the upper, rigid crust and 2) the earthquakes tend to concentrate above the discontinuities unveiled by the seismic exploration in the deep crust and at the Moho boundary. In this paper a similar analysis is conducted, even in volumes where DSS information is not available, with the goal of shedding some light on the continuation of these structures with depth. It is apparent that the upper mantle seismicity is very unevenly distributed; therefore, we only focus on the areas where a sub-crustal seismicity is recorded, adding to the seismic models of the crust some information, if available, on the physical characters of the upper lithosphere. Four areas are examined: the well-known Calabrian (Aeolian) Arc where the Ionian plate is subducted beneath the Tyrrhenian, thin crust of oceanic type, the active subduction of the slab being witnessed by deep and very deep earthquakes; the north-central Apennines where the continental crust of the Adria microplate seems also subducted beneath the transitional, peri-Tyrrhenian type of crust but where the observed hypocenters are limited to the depth of about 100 km; the northern Apennines, where the same type of subduction seems to occur beneath the north-eastern slope of the mountain range, though evidenced by an even smaller number of events; finally, the western Alps: also here a small group of foci are recorded in the upper mantle beneath the southern end of the "Ivrea body". The different behavior of deep seismicity in the four areas confirms that the Italian peninsula is formed by sectors deriving from different geodynamical processes.

### 1. Foreword

In a preceding paper (Cassinis and Solarino, 2006), the results of an analysis investigating the relationship between seismicity and the main features of the crust beneath the Italian peninsula were presented. In that study, two data sets were compared: the interpretation of the Deep Seismic Soundings - (DSS) along several transects, crisscrossing the Alpine range and the Italian peninsula (Cassinis *et al.*, 2003, 2005), and an updated catalog of the earthquakes (Chiarabba *et al.*, 2005). The most important conclusion was that, while the majority of the earthquakes happens in the rigid, upper crust, the hypocenters are assembled above the discontinuities

discovered by the DSS in the lower crust and at the Moho boundary. Since the maximum depth of the DSS is limited to the crust-mantle boundary, little information was obtained on the continuation with depth of the tectonic structures below that limit. Therefore, the upper bend of the subduction zones was imaged and investigated only in few cases.

Royden *et al.* (1987) had already pointed out that subduction-zone processes and their control of surface deformation are poorly understood, largely because few direct methods are available for observation of such a process. Nevertheless, some steps towards a more detailed knowledge of the subducting plates have been achieved in the last years using indirect techniques, like, for example, seismic tomography [see for example Solarino *et al.* (1996) or Di Stefano *et al.* (1999)]. The increasing number of available images of the subduction areas have favored several hypotheses on their nature and evolution; a discussion of these models is beyond the scope of this paper and will not be treated here. In this study, we integrate the information on the crust, already described by the seismic exploration, with a very well constrained distribution of seismicity, in order to infer the continuation at depth of the most debated structures.

In Fig. 1, the hypocenters recorded in the Italian region during the period 1981-2002 (Chiarabba *et al.*, 2005) are projected on a NW-SE cross-section; it is clear that they are concentrated within a depth of 25 - 30 km; however, five areas, where a sub - crustal seismicity is recorded, are observed namely, from NW to SE, the southwestern Alps, the northern Apennines, the north-central Apennines, the southeastern Alps (northern Dinarides) and, finally, Calabria-south Tyrrhenian Sea (the well-known Aeolian slab). From this synthetic approach, it can be remarked that the maximum depth reached by the earthquakes, as well as the number of events in the areas above are increasing while proceeding from NW to SE.

In Fig. 2, the distribution of magnitude versus depth is plotted for the part of the database where this parameter is known (about 50% of the events); it is shown that the threshold of the high magnitude events ( $> 5$ ) is found within a depth of about 30 km.

## 2. Seismic activity and crustal domains

In Fig. 3, the crustal domains, the Moho depth contour lines and the tectonic features in the deep crust and at the crust-mantle boundary are described according to a revision of the interpretation of DSS lines (Cassinis *et al.*, 2003, 2005; Cassinis and Solarino, 2006). The traces of the interpretative transects to be discussed later, are plotted: the empty ones indicate the cross-sections along which the interpretation of the crustal structure is available, while the solid grey traces correspond to where only the seismic events are plotted. In both cases, the hypocenters contained in a variable width (20 to 30 km) are projected on the vertical section.

In Fig. 4, the epicenters are plotted on the map of Fig. 3. Two classes of depth are considered:  $Z < 35$  km (mainly inside the crust; Fig. 4a), and  $Z > 35$  km (mainly in the upper lithosphere; Fig. 4b). Such a value was chosen on the basis of the average position of the Moho for the peninsula (Fig. 3) and on the findings of Chiarabba *et al.* (2005), who underlined a very clear difference in the seismicity above and below that depth.

The magnitude is also shown. The accuracy of both vertical and horizontal coordinates is within  $\pm 3$  km for all events: it allows us to discard the events with an arbitrarily depth assigned during the location process; furthermore, in this way, the uncertainty in the hypocentral position is




Fig. 1 - Summary of the seismicity in the Italian region according to the catalog by Chiarabba *et al.* (2005). All hypocenters recorded in the Italian region are projected on a NW-SE cross section. It is clear that the focal depth of the majority of the earthquakes is less than about 35 km. The foci slightly deepen while proceeding southwards. The areas where sub crustal events are observed are marked by numbers in the cross section: 1: western Alps, 2: north Apennines, 3: north-central Apennines, 4: SE Alps-Dinaric Alps, 5: Ionian-Aeolian slab.




Fig. 2 - Distribution of magnitude versus depth for the seismicity in the Italian region, showing that the majority of the high magnitude events ( $> 5$ ) is recorded within the depth of about 30 km. The group of deep events (around 200 km) with  $M > 5$  is located along the Aeolian slab.

comparable to the size and depth of the cross-sections traced on the map. For comments on the correlation of the epicenters position with the interpreted features in the crust and in the Moho boundary, reference is made to Cassinis and Solarino (2006). In comparing the two maps (Figs. 4a and 4b), the north westward shifting of the deep epicenters of the sinking Aeolian slab is very clear.

### 3. Description of seismic activity on the selected cross-sections

In the five investigated areas, a sub-crustal seismicity is recorded and reliable models of the crust are available. For the sake of clarity, the DSS cross-sections are assigned the same numbers that were used in the previous papers (Cassinis *et al.*, 2003, 2005; Cassinis and Solarino, 2006).

Let's start examining the area of the Calabrian (Aeolian) Arc; in Fig. 5 the interpretation of the DSS transect 9 - 9 (Steinmetz *et al.*, 1983) from the center of the Tyrrhenian Sea to Calabria and to the Ionian Sea is proposed, and all the available hypocenters are projected on the vertical section; they clearly show the shape of the subducting slab, down to the depth of about 300 km. The onset of the slab is in the Moho step that marks the transition from the margin of the Ionian domain to the Tyrrhenian, thinner crust. Note that the crust of the Ionian plate (Makris *et al.*, 1986; Scarascia *et al.*,




Fig. 3 - Depth contour lines of the Moho boundary (contour interval 2.5 km) and crustal domains modified [from Cassinis *et al.* (2003)]; the position of the cross-sections described in this paper is also shown.

The empty rectangles represent the transects where the interpretation of the seismic structure is available [the numbers are the same as those used in Cassinis *et al.* (2003)] while the solid grey rectangles correspond to the cross-sections where only the earthquake hypocenters are shown. The foci contained in a volume of variable width (from 20 to 30 km) are projected on each cross-section.

Explanation of symbols:

Crustal types: 1: European plate; 2: Afro-Adriatic plate; 3: Styrian and Pannonian basins; 4: Ligurian, Tuscan-Perityrrhenian transitional crust, the same ornamentation is used for the Pantelleria rift (Sicily Channel); 5: oceanic-sub-oceanic crust; 6: over-thrusting fronts of the Moho boundary of the Adriatic over the European plate (Alpine range) of the Ligurian, Tuscan, Perityrrhenian transitional crust over the Adriatic-African plate (Apennines range), of the Ligurian-Tuscan over the European (Corsica); 7: fragmentation lines in the upper mantle; 8: Moho depth contour lines (km); 9: Moho depth contour-lines (subducted).




Fig. 4 - Epicenters of the earthquakes according to the Chiarabba *et al.* (2005) catalog, plotted on the map in Fig. 3: a) focal depth  $Z < 35$  km; b) focal depth  $Z > 35$  km. Classes of magnitude are also indicated; white crosses are used for events whose magnitude is not computed.




Fig. 5 - Interpretation of the DSS profile 9- 9 crossing the Tyrrhenian Sea, mainland Calabria down to the Ionian Sea. Numerical values indicate the velocities of P waves in km/s. The hypocenters, both in the crust and in the upper mantle, show the shape of the Ionian slab that is subducted north-westwards beneath the Tyrrhenian Sea; the onset of the subduction is clearly situated in the Moho step that marks the transition from the margin of the Ionian domain to the Tyrrhenian oceanic crust.

1994) is thicker (about 20 km instead of 10 - 15 km of the Tyrrhenian) and is characterized by a thick sedimentary, low velocity cover.

In Fig. 6, cross-section 11a is described from the Aeolian Islands to Calabria, the Gulf of Taranto and reaching the tip of the Salentina Peninsula. Note the contrast between the SW part of the section (strong seismic activity beneath the Aeolian Islands and Calabria where the beginning of the slab is observed) and the NE portion (continental type of crust belonging to the African plate) where the earthquakes are almost absent.

The seismic foci are also plotted on three cross-sections directed nearly normally to section 11a; the north-eastern section shows the termination of the slab in that direction. While in section 11a, the seismic activity is continuous, though of variable magnitude along the whole slab, in the two cross-sections traced south-westwards two gaps, respectively at the depth of about 130 km and 60 km, can be observed.

The second investigated area is in the north-central Apennines; Fig. 7a illustrates the interpretation of cross-section 7-7 from Corsica to the Elba Channel, south Tuscany, the Apenninic range, the city of Perugia, the Tiber valley and the Adriatic coast near the city of Ancona. The highest




Fig. 6 - a) Cross-section 11a from the Aeolian Islands to Calabria, the Gulf of Taranto and to the tip of the Salentina Peninsula. Note the contrast between the south-western part of the section (strong seismic activity beneath the Aeolian Islands and Calabria (beginning of the slab) and the northeastern portion (continental type of crust belonging to the African plate) where the earthquakes are almost absent.

b) Cross-sections directed perpendicularly towards section 11a, showing the lateral continuation of the Aeolian slab. Note the gap in the slab at a depth of about 130 km (central section) and a minor gap at about 60 km in the south-western section.






Fig. 7 - a) Cross-section 7-7 from Corsica to the Elba Channel, southern Tuscany, north-central Apennines, Tiber Valley and the Adriatic coast near the city of Ancona, interpreted after the DSS data. The highest magnitude seismicity is concentrated in the upper crust, above the step of the Moho boundary near the city of Perugia (eastern side of the Apennines). The foci deepening westwards beneath the Adriatic coast seem to indicate the beginning of the subduction of the Adria microplate; the seismicity appears limited to a depth of about 80 - 90 km .  
 b) A line parallel to section 7-7 (about 80 km south) showing that the deep earthquakes have almost disappeared.

magnitude seismicity is concentrated in the upper crust, above the step of the Moho boundary near Perugia (eastern side of the Apennines).

The foci (maximum depth 80-90 km) deepening westwards beneath the Adriatic coast seem to indicate the beginning of the subduction of the Adria microplate. Here the trend of the hypocenters agrees with the interpretation of the deep reflection line CROP 03 that reaches the coast at Pesaro, about 60 km north of Ancona; the deeper recorded events (mainly diffractions) outline the westward immersion of the thicker Adria crust, in contrast with the flat pattern peculiar of the Tuscan-Tyrrhenian thin crust, characterized by high heat flow [see the interpretative models by Barchi *et al.* (1998), Decandia *et al.* (1998), Gualteri *et al.* (1998), Cassinis (2002) and also Cassinis and Solarino (2006)]. In Fig 7b, a line, parallel to section 7-7 (about 80 km south), shows that the deep earthquakes have almost disappeared.

In Fig 8, the DSS line 6-6 is illustrated (see also Letz *et al.*, 1977; Scarascia *et al.*, 1994), from the Ligurian Sea, north of Corsica, to the coast of northern Tuscany near Viareggio, then crossing the

northern Apennines. A moderate seismic activity is observed beneath the mountain range and also here, a small group of deep foci (70 - 80 km) is shown. The foci are plotted on sections perpendicular to line 6-6: no seismicity in the upper mantle is recorded beneath the coast of northern Tuscany and the Ligurian Sea.

Tomographic images, obtained by using different methods, show a pronounced high-velocity anomaly in the northern Apennines, interpreted as a submerged crustal slab, representing the westward continuation of the Adriatic Moho. The shape and location of the imaged, high velocity slab are slightly different, reflecting a difference in the model resolution obtained by using local, regional and teleseismic data. In particular, the slab continuity is still debated. Lucente *et al.* (1999) and Piromallo and Morelli (2003), define a continuous slab down to a 670 km depth, while Spakman and Wortel (2004), hypothesize a relatively short (300-400 km) and continuous northern Apennines slab which may strongly curve to the west beneath the Po plain. In their view, a short north-Apennines slab is sufficient to explain the opening history of the northern Liguro-Provençal basin. Moreover, the negative anomalies imaged in their tomographic images to depths of 200 km under the central-southern Apennines are attributed to slab detachment. Such geometry would justify the absence of earthquakes beneath 100 km under the whole central Apennines.

The cross-sections across the western Alps are illustrated in Fig. 9. The interpretation of the central section (3), [Grenoble-Turin-Po Valley; for an earlier recorded one, see also Giese and Prodehl (1976)] is controversial; the one presented here (Cassinis *et al.*, 2003, 2005) was assumed as the most likely: the shallow fragment, having a velocity of about 7.5 km/s, is attributed to the lifted and impoverished mantle material belonging to the Adriatic-African domain, overthrusting the European Moho that is subducted south-westwards, beneath the Po Valley. The deep reflection line (Nicolas *et al.*, 1990) did not confirm this hypothesis, that is, in turn, supported by gravity anomalies. Along the section the seismicity appears to be confined to the upper crust, especially in the western side of the Alps; the foci very clearly follow the shape of the Moho fragment (“Ivrea body”) on the eastern side. A similar pattern is observed along the parallel sections except in the southernmost one, where a small group of subcrustal events is recorded, positioned in the area where the European, Adria and Ligurian domains meet. The results of passive seismic tomography (Solarino *et al.*, 1997; Eva *et al.*, 2001; Scafidi *et al.*, 2006) show that an anomalous high velocity is found in the same area, in the upper mantle. This location corresponds also to the sudden change of direction of the gravity anomalies [see the insert in Fig. 9, Klingele *et al.* (1992)]. Therefore, a different geodynamic behaviour of the sectors of the Alpine range is proposed and evidenced by the comparison of the two data sets used in this work.

#### 4. Concluding remarks

The main conclusion of a previous paper (Cassinis and Solarino, 2006) was that the seismic regime in the Italian region changes according to the crustal structure and the Moho boundary characteristics.

In this paper, particular attention is given to the sub-crustal seismicity. It has to be remarked that the joint interpretation of DSS and seismicity alone cannot provide a complete frame of the structural, deep setting of the Italian peninsula. This happens for several reasons: the discreteness of information, the inhomogeneous distribution of seismic stations, with consistent gaps especially in




Fig. 8 - a) DSS line 6-6 from the Ligurian Sea, north of Corsica, to the coast of northern Tuscany near Viareggio, then crossing the northern Apennines. A moderate seismic activity is observed beneath the mountain range and a small group of deep foci (up to 80 km deep) is also shown. b) The foci are plotted on sections perpendicular to line 6-6: no seismicity in the upper mantle is recorded beneath the Ligurian Sea.




Fig. 9 - Interpretative transects across the western Alps. The central line 3-3 (from Grenoble to Turin and the Po Valley) illustrates the crustal structure interpreted mainly after the DSS results. The interpretation of passive tomography in the crust and in the upper mantle (after Scafidi *et al.*, 2006) is also shown in this section as well as in parallel sections (see the colour scale of P velocity in km/s). Note the group of sub-crustal events along the southernmost cross-section. Further comments in the text. The traces of cross-sections are also plotted (see the enclosed map) on the Bouguer gravity anomalies (Klingelé *et al.*, 1992).

the seas, the threshold of magnitude, and many others. Furthermore, it must be reminded that the maximum penetration of DSS is of about 60-70 km. Nevertheless, the analysis highlights the complexity of the geological structure and the different geodynamic history of each sector of the Alpine range, of the Apennines as well as of the Calabrian Arc and Sicily. Earthquakes in the upper mantle are observed especially where a process of subduction is taking place. However, such process occurs with different mechanisms.

The following synthesis of the results can be made.

- The lateral extension of the well known Aeolian-Calabrian-Ionian slab appears well defined, corresponding to the north-eastern edge of the Ionian plate. The relationships between the crust and the upper mantle are clear: the onset of the subducting slab is in the step marking the transition from the Tyrrhenian (thin, oceanic type of crust) to the Ionian (thicker, oceanic type of crust). Unfortunately, in the Ionian plate, the knowledge of some complementary data (heat flow, seismic activity) is not adequate for a more quantitative interpretation. Nevertheless, the geometry, size and characteristics of the slab have been extensively studied by many tomographic experiments. Di Stefano *et al.* (1999) have underlined the presence of two broad low-velocity zones located beneath both the south-eastern portion of the Calabrian arc and the Aeolian Islands, at slightly different depths. Our results show that, while the rigid behaviour of the main part of the subducting material is witnessed by the continuity of the earthquakes along the sinking slab down to about 300 km (cross-section 9-9, Fig. 5), this may be questioned when looking at Fig. 6. In fact, while section 11a (top central panel) clearly exhibits a continuous slab, the perpendicular sections show a gap of seismicity at different depths. Moreover, it is evident that the higher magnitude events are the deepest ones. A possible explanation for this remark is that the slab is continuous but has different rheological properties within its extension, as the tomography seems to suggest. Where the velocity is lower (due to petrological or thermal factors), seismicity is almost absent or at least less frequent. The magnitudes are also reduced, being the slab in a non perfectly-fragile condition in that portion.
- A completely different characteristic is shown in the two investigated areas along the Apennines (north-central and northern). In both areas, the depth and number of sub-crustal earthquakes are comparable, the seismic activity being slightly higher in the former. The structure of the crust is also similar, the continental, thicker Adriatic-Padan crust (low heat flow) subducting beneath the transitional, thin peri-Tyrrhenian (high heat flow) type of crust (Mongelli *et al.*, 1991). The sub-crustal earthquakes are not deeper than about 100 km; the deep foci (Fig. 4b) tend to concentrate beneath the bowed, broken line marking the edge of the over thrusting peri-Tyrrhenian Moho.

The northernmost investigated area is situated near the southern end of the western Alps (Figs. 3 and 4b), close to the observed inversion of the over-thrusting of the Moho boundary: in the Alpine range the external plate (European) is subducted beneath the Adria plate while, more eastwards, the latter domain is subducted beneath the peri-Tyrrhenian Moho. In this area, a small group of sub-crustal earthquakes is positioned beneath this area of inversion. The correct position of these events, can be considered as not questionable, since Cattaneo *et al.* (1999) obtained a similar result analyzing the seismicity of the south-western Alps using a much denser seismic array and performing several location reliability tests. These events, no deeper than 60-70 km, even though according to Cattaneo *et al.* (1999), they may reach a depth of 120 km,

seem to be related to high velocity bodies (Scafidi *et al.*, 2006) that are located in the Monferrato area. Several authors have investigated the nature of these anomalous rocks, considering them a part of a trapped mantle (Schmid and Kissling, 2000) or a portion of an uplifted lower crust. These foci are very close to the region where the Bouguer anomalies (Klingelé *et al.*, 1992) show an abrupt change and where a sudden variation of the Moho depth (see Figs. 1 and 9) is supposed to be located.

A meaningful complement to seismicity would derive from the analysis of focal mechanisms. However, this would call for a selection (on the basis, for example, of the magnitude of seismic events) of the data for display among the many fault plane solutions available for the peninsula, obtained either by a first onset technique or a waveform inversion. To avoid this arbitrary selection, we believe that a sort of cumulative representation best suits the investigation in large areas. In this sense, a useful aid is given by the comprehensive survey by Vannucci *et al.* (2004): they display a map representing the seismic deformation obtained by a moment tensor summation on a 0.5 degree per 0.5 degree. The narrow and continuous belt of extensional deformation runs from the Strait of Messina to the Irpinia region and is also outlined by the extensional T-axes that are always perpendicular to the chain. The pattern of seismic deformation is less clear and more heterogeneous at the Gargano promontory latitude and northwards, being mainly characterized by compressive deformation in the outer part of the chain and in the foredeep. The transition between the southern and the central Apennines is characterized by compressional and strike-slip focal mechanisms. However, extension, perpendicular to the mountain belt continues all the way to the northern Apennines, as indicated by individual focal mechanisms and by P-axes of moment tensor sums.

To summarize, lateral heterogeneities, different thicknesses and a strongly irregular shape of the subducted Ionian-Adriatic lithosphere have produced a complex system of subduction in the Italian region. The two main arcs are characterized by the subduction of the oceanic lithosphere (Calabrian Arc) and of the continental lithosphere (north-central Apennines). In the area between the two arcs, the subduction process seems interrupted (Di Stefano *et al.*, 1999; Spakman and Wortel, 2004); the distribution of focal mechanisms, discussed above, seems to strengthen this hypothesis. Also to be remarked is that in the southern Apennines, where the Moho overthrusting appears to be not bowed but tends to line up, no earthquakes are recorded in the upper mantle (Fig. 4b). This could also support (Spakman and Wortel, 2004), the hypothesis of a progressive detachment north of the Calabrian slab.

**ACKNOWLEDGMENTS.** We are grateful to Claudio Eva and Roberto Sabadini for encouraging us to publish this paper and for giving suggestions that greatly improved the manuscript.

## REFERENCES

- Barchi M., Minelli G. and Piali G.; 1998: *The CROP 03 profile: a synthesis of results on deep structures of the Northern Apennines*. Mem. Soc. Geol. It., **52**, 383-400.
- Cassinis R.; 2002: *The Contribution of geophysical methods in integrated projects directed by Giampaolo Pialli*. Boll. Soc. Geol. It., Vol. Spec. n. **1**, 3-12.
- Cassinis R. and Solarino S.; 2006: *Seismicity and crustal structure in the Italian region: a new review using a synthesis of DSS results and updated catalogues of earthquakes*. Boll. Geof. Teor. Appl., **47**, 481-496.
- Cassinis R., Scarascia S. and Lozej A.; 2005: *The deep crustal structure of Italy and surrounding area from seismic refraction data. A new synthesis*. Boll. Soc. Geol. It., **122**, 365-376.
- Cassinis R., Scarascia S. and Lozej A.; 2005: *Review of seismic wide angle reflection-refraction (WARR) results in the Italian region (1956-1987)*. In: Finetti I.R. (ed), CROP Project - deep seismic exploration of the central Mediterranean region and Italy, Atlases in Geoscience, Vol. 1, Elsevier Earth and Environmental Science, Amsterdam, pp. 31-55.
- Cattaneo M., Augliera P., Parolai S. and Spallarossa D.; 1999: *Anomalous deep earthquakes in northwestern Italy*. J. of Seism., **3**, 421-435.
- Chiarabba C., Jovane L. and Di Stefano R.; 2005: *A new view of Italian seismicity using 20 years of instrumental recordings*. Tectonophysics, **395**, 251-268.
- Decandia F. A., Lazzarotto A., Liotta D. Cernobori L. and Nicolich R.; 1998: *The CROP 03 traverse: insights on post-collisional evolution of Northern Apennines*. Mem. Soc. Geol. It., **52**, 427-440.
- Di Stefano R., Chiarabba C., Lucente F. and Amato A.; 1999: *Crustal and uppermost mantle structure in Italy from inversion of P wave arrival times: geodynamic implications*. Geoph. Journ. Int., **139**, 483-498.
- Eva E., Solarino S. and Spallarossa D.; 2001: *Seismotectonics and local earthquake tomography of the western Ligurian Sea*. Tectonophysics, **339**, 391-406.
- Giese P., and Prodehl, C.; 1976: *Main features of crustal structures of the Alps*. In: Giese P., Prodehl C. and Stein A. (eds), Explosion seismology in central Europe, Deutsch. Geophys. Gesell., Springer, pp. 347-375.
- Gualteri L., Augliera P., Eva C., and Cassinis R.; 1998: *Integration of active and passive seismic investigations: constraints on the geodynamics of the northern Apennines*. Mem. Soc. Geol. It., **52**, 327-336.
- Klingelé E., Lahmeyer B. and Freeman F.; 1992: *Bouguer gravity anomalies*. In: Freeman R. and Mueller S. (eds), The European Geotraverse, A continent revealed. Atlas of compiled data, pp. 27-31.
- Letz H., Reichert C. and Wigger P.; 1977: *Results of two seismic refraction lines in the Northern Apennines (lines 1 and 3)*. Boll. Geof. Teor. Appl., **19**, 225-232.
- Lucente F. P., Chiarabba C., Cimmini, G. B. and Giardini, D.; 1999: *Tomographic constraints on the geodynamic evolution of the Italian region*. J. Geophys. Res., **104**, 20307-20327.
- Makris J., Nicolich R. and Weigel C.; 1986: *A seismic study in the Western Ionian sea*. Annales Geophysicae, **4**, 665-678.
- Mongelli F., Zito G., Della Vedova B., Pellis G., Squarci P. and Taffi L.; 1991: *Geothermal regime of Italy and surrounding seas*. In: Cermak V. and Rybach L. (eds), Exploration of the deep continental crust, Springer-Verlag, Berlin.
- Nicolas A., Polino R., Hirn A., Nicolich R. and ECORS-CROP W.G.; 1990: *ECORS-CROP traverse and deep structure of the western Alps: a synthesis*. In: Roure F. and Polino R. (eds), Deep structure of the Alps, Vol. Spec. Soc. Geol. It., Roma, pp. 15-27.
- Piomallo, C. and Morelli, A.; 2003: *P-wave tomography of the top 1000 km under the Alpine-Mediterranean area*. J. Geophys. Res., **108**, doi:10.1029/2002JB001757.
- Royden L., Patacca E. and Scandone P.; 1987: *Segmentation and configuration of subducted lithosphere in Italy: An important control on thrust-belt and foredeep-basin evolution*. Geology, **15**, 714-717.
- Scafidi D., Solarino S. and Eva C.; 2006: *Structure and properties of the Ivrea body and the Alps-Apennines system as revealed by local earthquake tomography*. Boll. Geof. Teor. Appl., **47**, 497-514.
- Scarascia S., Lozej A. and Cassinis R.; 1994: *Crustal structures of the Ligurian, Tyrrhenian and Ionian seas and adjacent onshore areas interpreted from wide angle seismic profiles*. Boll. Geof. Teor. Appl., **35**, 5-21.
- Schmid S.M. and Kissling E.; 2000: *The arc of the western Alps in the light of geophysical data on deep crustal structure*. Tectonics, **19**, 62-85.
- Solarino S., Spallarossa D., Parolai S., Cattaneo M. and Eva C.; 1996: *Lithoasthenospheric structures of Northern Italy as inferred by teleseismic waves tomography*. Tectonophysics, **260**, 271-289.

- Solarino, S., Kissling, E., Sellami, S., Smriglio, G., Thouvenot, F., Granet, M., Bonjer, K.P. and Slejko D.; 1997: *Compilation of a recent seismicity data base of the greater Alpine region from several seismological networks and preliminary 3D tomographic results*. *Annali di Geofisica*, **50**, 61-174.
- Spakman W. and Wortel R.; 2004: *A Tomographic View on Western Mediterranean Geodynamics*. In: Cavazza W., Roure F., Spakman W., Stampfli G.M. and Ziegler P.A. (eds), *The TRANSMED Atlas. The Mediterranean Region from Crust to Mantle. Geological and Geophysical Framework*, Springer.
- Steinmetz L., Ferrucci F., Hirn A., Morelli C. and Nicolich R. ; 1983: *A 550 km long Moho traverse in the Tyrrhenian Sea from OBS recorded Pn waves*. *Geophys. Res. Letters*, **10**, 428-431.
- Vannucci G. and Gasperini P.; 2004: *The new release of the database of earthquake mechanisms of the Mediterranean area (EMMA version 2)*. *Annals of Geophysics*, **47**, 307-326.

*Corresponding author:* Stefano Solarino  
Istituto Nazionale di Geofisica e Vulcanologia, CNT  
c/o Dipteris, Università di Genova  
Viale Benedetto XV, 5, 16132 Genova, Italy  
e-mail: solarino@ingv.it